

## STEPENOVARANJE

Proizvod  $a \cdot a \cdot \dots \cdot a = a^n$  naziva se  $n$ -tim stepenom broja. Ako je  $a \in R$ ,  $a \neq 0$  i neka je  $n \in N$   
 $\underset{n\text{-puta}}{\dots}$

Po definiciji je:

$$\begin{aligned} \mathbf{1)} \quad a^0 &= 1 \rightarrow \text{primer: } 5^0 = 1, \ (-3)^0 = 1, \ \left(\frac{4}{7}\right)^0 = 1 \\ \mathbf{2)} \quad a^{-n} &= \frac{1}{a^n} \rightarrow \text{primer: } 3^{-2} = \frac{1}{3^2} = \frac{1}{9}, \ 5^{-3} = \frac{1}{5^3} = \frac{1}{125} \end{aligned}$$


---

Još važe sledeća pravila:

$$\begin{aligned} \mathbf{3)} \quad a^m \cdot a^n &= a^{m+n} \quad \rightarrow \text{primer: } 3^2 \cdot 3^5 = 3^{2+5} = 3^7 \\ \mathbf{4)} \quad a^m : a^n &= a^{m-n} \quad \rightarrow \text{primer: } 7^{10} : 7^6 = 7^{10-6} = 7^4 \\ \mathbf{5)} \quad (a^m)^n &= a^{m \cdot n} \quad \rightarrow \text{primer: } (2^3)^5 = 2^{3 \cdot 5} = 2^{15} \\ \mathbf{6)} \quad (a \cdot b) &= a^n \cdot b^n \quad \rightarrow \text{primer: } (12 \cdot 11)^5 = 12^5 \cdot 11^5 \\ \mathbf{7)} \quad \left(\frac{a}{b}\right)^n &= \frac{a^n}{b^n} \quad \rightarrow \text{primer: } \left(\frac{7}{4}\right)^2 = \frac{7^2}{4^2} \\ \mathbf{8)} \quad \left(\frac{a}{b}\right)^{-n} &= \left(\frac{b}{a}\right)^n \quad \rightarrow \text{primer: } \left(\frac{2}{3}\right)^{-2} = \left(\frac{3}{2}\right)^2 = \frac{3^2}{2^2} = \frac{9}{4} \end{aligned}$$

O čemu treba voditi računa?

Treba paziti na zapis:  $(-5)^2 = (-5)(-5) = 25$ , dok  $-5^2 = -5 \cdot 5 = -25$ . Uopšteno važi:

$$\begin{aligned} (-a)^{\text{paran}} &= a^{\text{paran}} \\ (-a)^{\text{neparan}} &= -a^{\text{neparan}} \end{aligned}$$

Dakle, paran izložilac “uništiti” minus.

## ZADACI

1) Izračunati:  $\frac{(2^7 : 2^5) \cdot 2^3}{2^4 : 2^2}$

$$\frac{(2^7 : 2^5) \cdot 2^3}{2^4 : 2^2} = \frac{2^{7-5} \cdot 2^3}{2^{4-2}} = \frac{2^2 \cdot 2^3}{2^2} = \frac{2^{2+3}}{2^2} = \frac{2^5}{2^2} = 2^{5-2} = 2^3 = 8$$

---

2) Izračunati:  $\frac{3^5 \cdot 9^3}{27^2 \cdot 3}$

$$\frac{3^5 \cdot 9^3}{27^2 \cdot 3} = \frac{3^5 \cdot (3^2)^3}{(3^3)^2 \cdot 3^1} = \frac{3^5 \cdot \cancel{3^6}}{\cancel{3^6} \cdot 3^1} = \frac{3^5}{3^1} = 3^{5-1} = 3^4 = 3 \cdot 3 \cdot 3 \cdot 3 = 81$$

---

3) Izračunati:  $\frac{(x^4)^3 \cdot x^3 : x^5}{(x^5 : x^2)^3} =$

$$\frac{(x^4)^3 \cdot x^3 : x^5}{(x^5 : x^2)^3} = \frac{x^{12} \cdot x^3 : x^5}{(x^{5-2})^3} = \frac{x^{12+3-5}}{(x^3)^3} = \frac{x^{10}}{x^9} = x^{10-9} = x^1 = x$$

---

4) Izračunati:  $\frac{3^{n+1} \cdot 3^{n+2}}{3^{2n+4}}$

$$\begin{aligned} \frac{3^{n+1} \cdot 3^{n+2}}{3^{2n+4}} &= \frac{3^{n+1+n+2}}{3^{2n+4}} = \frac{3^{2n+3}}{3^{2n+4}} = \text{Pazi pa zagrade zbog minusa} \\ &= 3^{(2n+3)-(2n+4)} = 3^{2n+3-2n-4} = 3^{-1} = \frac{1}{3^1} = \frac{1}{3} \end{aligned}$$

---

5) Izračunati  $0,5^{-1} + 0,25^{-2} + 0,125^{-3} + 0,0625^{-4}$

$$0,5^{-1} + 0,25^{-2} + 0,125^{-3} + 0,0625^{-4} =$$

$$\left(\frac{1}{2}\right)^{-1} + \left(\frac{1}{4}\right)^{-2} + \left(\frac{1}{8}\right)^{-3} + \left(\frac{1}{16}\right)^{-4} =$$

$$\left(\frac{2}{1}\right)^1 + \left(\frac{4}{2}\right)^2 + \left(\frac{8}{1}\right)^3 + \left(\frac{16}{1}\right)^4 =$$

$$2^1 + 4^2 + 8^3 + 16^4 = 2 + 16 + 512 + 65536 = 66066$$


---

6) Izračunati  $1^{-1} + 2^{-2} + 3^{-3} + (-1)^{-1} + (-2)^{-2} + (-3)^{-3}$

$$1^{-1} + 2^{-2} + 3^{-3} + (-1)^{-1} + (-2)^{-2} + (-3)^{-3} =$$

$$\frac{1}{1} + \frac{1}{2^2} + \frac{1}{3^3} + \frac{1}{(-1)^1} + \frac{1}{(-2)^2} + \frac{1}{(-3)^3} =$$

$$1 + \frac{1}{4} + \frac{1}{27} - 1 + \frac{1}{4} - \frac{1}{27} = \frac{1}{4} + \frac{1}{4} = \frac{2}{4} = \frac{1}{2}$$


---

7) Ako je  $a = 5^3 \cdot \left(\frac{1}{4}\right)^{-4} \cdot \left(\frac{3}{2}\right)^2$  i  $b = 10^3 \left(\frac{5}{3}\right)^{-2}$  nadji  $a \cdot b^{-1}$

$$a = 5^3 \cdot \left(\frac{1}{4}\right)^{-4} \cdot \left(\frac{3}{2}\right)^2 = 5^3 \cdot \left(\frac{4}{1}\right)^4 \cdot \frac{3^2}{2^2} = \frac{5^3 \cdot 4^4 \cdot 3^2}{2^2}$$

$$= \frac{5^3 \cdot (2^2)^4 \cdot 3^2}{2^2} = 5^3 \cdot (2^2)^3 \cdot 3^2 = 5^3 \cdot 2^6 \cdot 3^2$$

$$b = 10^3 \cdot \left(\frac{5}{3}\right)^{-2} = 10^3 \cdot \left(\frac{3}{5}\right)^2 = \frac{10^3 \cdot 3^2}{5^2} = \frac{(5 \cdot 2)^3 \cdot 3^2}{5^2} = \frac{5^3 \cdot 2^3 \cdot 3^2}{5^2} = 5 \cdot 2^3 \cdot 3^2$$

Konačno: izračunati i  $a \cdot b^{-1}$

$$a \cdot b^{-1} = 5^3 \cdot 2^6 \cdot 3^2 \cdot \frac{1}{5 \cdot 2^3 \cdot 3^2} = 5^2 \cdot 2^3 = 25 \cdot 8 = 200$$


---

8) Izračunati  $\left( \left( \frac{5x^{-5}}{2y^{-2}} \right)^{-2} \cdot \left( \frac{y^{-1}}{5x^{-1}} \right)^{-3} \right) : 10x^2y^{-3}$

$$\left( \left( \frac{5x^{-5}}{2y^{-2}} \right)^{-2} \cdot \left( \frac{y^{-1}}{5x^{-1}} \right)^{-3} \right) : 10x^2y^{-3} =$$

$$\left( \frac{5^{-2} \cdot x^{10}}{2^{-2} \cdot y^4} \cdot \frac{y^3}{5^{-3} \cdot x^3} \right) : 10x^2y^{-3} =$$

$$(5^{-2+3} \cdot x^{10-3} \cdot y^{3-4} \cdot 2^2) : 10x^2y^{-3} =$$

$$(5^1 \cdot x^7 \cdot y^{-1} \cdot 4) : 10x^2y^{-3} =$$

$$\frac{20}{10} x^{7-2} y^{-1-(-3)} = 2x^5 y^{-1+3} = 2x^5 y^2$$


---

9) Ako je  $10^x = \frac{\frac{1}{2}10^{-3} + \frac{1}{2}10^{-4}}{55 \cdot 10^{-7}}$  Odrediti x.

$$\frac{\frac{1}{2}10^{-3} + \frac{1}{2}10^{-4}}{55 \cdot 10^{-7}} = \frac{\frac{1}{2000} + \frac{1}{20000}}{\frac{55}{10000000}} = \text{Izvučemo gore zajednički}$$

$$\frac{\frac{1}{2000} \left( 1 + \frac{1}{10} \right)}{\frac{55}{10000000}} = \frac{\frac{1}{2000} \cdot \frac{11}{10}}{\frac{55}{10000000}} = \frac{\frac{11}{20000}}{\frac{55}{10000000}} =$$

$$\frac{11 \cdot 10000000}{20000 \cdot 55} = \frac{11 \cdot 10000000}{11 \cdot 100000} = \frac{10000000}{100000} = 100 = 10^2$$

sada je  $10^x = 10^2$ , dakle  $x = 2$

---

10)

a)  $A \cdot 10^{-5} = 0,2 \cdot 0,008$

$$A \cdot 10^{-5} = 2 \cdot 10^{-1} \cdot 8 \cdot 10^{-3}$$

$$A \cdot 10^{-5} = 16 \cdot 10^{-4}$$

$$A = \frac{16 \cdot 10^{-4}}{10^{-5}}$$

$$A = 16 \cdot 10^{-4-(-5)}$$

$$A = 16 \cdot 10^{-4+5}$$

$$A = 16 \cdot 10$$

$$A = 160$$

b)  $B \cdot 10^{-6} = 0,04 \cdot 0,006$

$$B \cdot 10^{-6} = 4 \cdot 10^{-2} \cdot 6 \cdot 10^{-3}$$

$$B \cdot 10^{-6} = 24 \cdot 10^{-5}$$

$$B = \frac{24 \cdot 10^{-5}}{10^{-6}}$$

$$B = 24 \cdot 10^{-5+6}$$

$$B = 24 \cdot 10$$

$$B = 240$$

Ovde smo koristili zapisivanje realnog broja u sistemu sa osnovnim 10. Ovo je dobra opcija kada je broj "glomazan".

Primeri:

- 1) Brzina svetlosti je približno  $c \approx 300000000 m/s$  a mi je "lakše" zapisujemo  $c \approx 3 \cdot 10^8 m/s$ ,  $10^8$ -znači da ima 8 nula iza jedinice!!!
- 2)  $\frac{1}{500000} = \frac{1}{5 \cdot 10^5} = \frac{1}{5} \cdot 10^{-5} = \frac{2}{10} \cdot 10^{-5} = 2 \cdot 10^{-1} \cdot 10^{-5} = 2 \cdot 10^{-6}$
- 3)  $0,000069 = 6,9 \cdot 10^{-5} \approx 7 \cdot 10^{-5}$
- 4) Površina zemlje je  $510083000 km^2$  ali mi zapisujemo  $\approx 5 \cdot 10^8 km^2$

**11) Izračunati**  $\left( \frac{3a^{-x}}{1-a^{-x}} - \frac{2a^{-x}}{1+a^{-x}} - \frac{a^x}{a^{2x}-1} \right) : \frac{a^{-x}}{a^x-a^{-x}}$

$$\begin{aligned} & \left( \frac{3a^{-x}}{1-a^{-x}} - \frac{2a^{-x}}{1+a^{-x}} - \frac{a^x}{a^{2x}-1} \right) : \frac{a^{-x}}{a^x-a^{-x}} = \\ & \left( \frac{\frac{3}{a^x}}{1-\frac{1}{a^x}} - \frac{\frac{2}{a^x}}{1+\frac{1}{a^x}} - \frac{a^x}{a^{2x}-1} \right) : \frac{\frac{1}{a^x}}{a^x-\frac{1}{a^x}} = \\ & \left( \frac{\frac{3}{a^x}}{\frac{a^x-1}{a^x}} - \frac{\frac{2}{a^x}}{\frac{a^x+1}{a^x}} - \frac{a^x}{a^{2x}-1} \right) : \frac{\frac{1}{a^x}}{\frac{a^{2x}-1}{a^x}} = \\ & \left( \frac{3}{a^x-1} - \frac{2}{a^x+1} - \frac{a^x}{(a^x-1)(a^x+1)} \right) : \frac{1}{a^{2x}-1} = \\ & \frac{3(a^x+1)-2(a^x-1)-a^x}{(a^x-1)(a^x+1)} \cdot \frac{a^{2x}-1}{1} = \\ & \frac{3a^x+3-2a^x+2-a^x}{(a^x-1)(a^x+1)} \cdot \frac{(a^x-1)(a^x+1)}{1} = \\ & = 3+2=5 \end{aligned}$$

12) Izračunati  $\left( \frac{x-x^{-2}}{x^{-2}+x^{-1}+1} - \frac{x-x^{-1}}{1+x^{-2}+2 \cdot x^{-1}} \right) : \frac{1-x^{-1}}{1+x^{-1}}$

$$\begin{aligned}
& \left( \frac{x-x^{-2}}{x^{-2}+x^{-1}+1} - \frac{x-x^{-1}}{1+x^{-2}+2 \cdot x^{-1}} \right) : \frac{1-x^{-1}}{1+x^{-1}} = \\
& \left( \frac{x-\frac{1}{x^2}}{\frac{1}{x^2}+\frac{1}{x}+1} - \frac{x-\frac{1}{x}}{1+\frac{1}{x^2}+\frac{2}{x}} \right) : \frac{1-\frac{1}{x}}{1+\frac{1}{x}} = \\
& \left( \frac{\frac{x^3-1}{x^2}}{\frac{1+x+x^2}{x^2}} - \frac{\frac{x^2-1}{x}}{\frac{x^2+1+2x}{x^2}} \right) : \frac{\frac{x-1}{x}}{\frac{x+1}{x}} = \\
& \left( \frac{(x-1)(x^2+x+1)}{\cancel{x^2+x+1}} - \frac{x(x-1)(x+1)}{(x+1)^2} \right) : \frac{x-1}{x+1} = \\
& \left( \frac{x-1}{1} - \frac{x(x-1)}{x+1} \right) : \frac{x-1}{x+1} = \\
& \frac{(x-1)(x+1)-x(x-1)}{x+1} \cdot \frac{x+1}{x-1} = \\
& \frac{\cancel{(x-1)}[(x+1)-x]}{\cancel{x+1}} \cdot \frac{\cancel{x+1}}{\cancel{x-1}} = x+1-x=1
\end{aligned}$$

**13) Izračunati**  $A = \left( \frac{a^n}{1-a^{-n}} + \frac{a^{-n}}{1+a^{-n}} \right) - \left( \frac{a^n}{1+a^{-n}} + \frac{a^{-n}}{1-a^{-n}} \right)$

$$A = \left( \frac{a^n}{1-a^{-n}} + \frac{a^{-n}}{1+a^{-n}} \right) - \left( \frac{a^n}{1+a^{-n}} + \frac{a^{-n}}{1-a^{-n}} \right)$$

$$A = \left( \frac{\frac{a^n}{1}}{\frac{1}{a^n}-1} + \frac{\frac{1}{a^n}}{\frac{1}{a^n}+1} \right) - \left( \frac{\frac{a^n}{1}}{\frac{1}{a^n}+1} + \frac{\frac{1}{a^n}}{\frac{1}{a^n}-1} \right)$$

$$A = \left( \frac{\frac{a^n}{1}}{\frac{a^n-1}{a^n}} + \frac{\frac{1}{a^n}}{\frac{a^n+1}{a^n}} \right) - \left( \frac{\frac{a^n}{1}}{\frac{a^n+1}{a^n}} + \frac{\frac{1}{a^n}}{\frac{a^n-1}{a^n}} \right)$$

$$A = \left( \frac{\frac{a^{2n}}{a^n-1}}{1} + \frac{1}{a^n+1} \right) - \left( \frac{\frac{a^{2n}}{a^n+1}}{1} + \frac{1}{a^n-1} \right)$$

$$A = \frac{a^{2n}(a^n+1)+1(a^n-1)}{(a^n-1)(a^n+1)} - \frac{a^{2n}(a^n-1)+1(a^n+1)}{(a^n+1)(a^n-1)}$$

$$A = \frac{a^{3n}+a^{2n}+a^n-1-(a^{3n}-a^{2n}+a^n+1)}{(a^n-1)(a^n+1)}$$

$$A = \frac{a^{3n}+a^{2n}+a^n-1-a^{3n}+a^{2n}-a^n-1}{(a^n-1)(a^n+1)}$$

$$A = \frac{2a^{2n}-2}{(a^n-1)(a^n+1)} = \frac{2(a^{2n}-1)}{(a^n-1)(a^n+1)} = \frac{2(a^n-1)(a^n+1)}{(a^n-1)(a^n+1)} = 2$$